Verde Valley Medical Center Off-Line Guidelines

2013 Changes (additional)

Off-line	Page	Change	Why
Adult Cardiopulmonary arrest- CCR alternative	27	Added a footnote to indicate escalating shocks for V-Fib/Pulseless V-Tach Footnote (6)	Per Medical control refractory V-Fib/ Pulseless V-Tach should have increasing joule dosing
Adult Pulseless Arrest	28	Added a footnote to indicate escalating shocks for V-Fib/Pulseless V-Tach Footnote (10)	Per Medical control refractory V-Fib/ Pulseless V-Tach should have increasing joule dosing
Adult Tachycardia with Pulses	30	Added a footnote requiring medical control input for use of amiodarone in patient with a systolic BP less than 90 mm HG	Amiodarone increases hypotension requiring medical control input.
Trauma-Burns	32	Added NS as option in fluid resuscitation	LR is still recommended solution for burns, included NS if LR unavailable
Pediatric Tachycardia with Pulses	62	Changed verbiage on footnote (6). Added contact medical control for use of Amiodarone if BP is less than 90 mm Hg systolic	Amiodarone increases hypotension requiring medical control input.
Pediatric Airway	74	Removed use of bougie for pediatrics with intubation attempts.	Current Bougie devices will only pass easily through a 6.0 ET tube.